REFERENCES

1. Agarwal A, (2005), Critical issues in Quality Control of Herbal Products, Pharma Times, vol 37(6):09-11.
2. Andrew Zusevics, (2008),Studies on winged bean, psophocarpus tetragonolobus (L.) nutrition in laboratory environment,Journal of plant nutrition.2008; volume 3.Issue 5.Page no.789-802.
3. Ankur R et.al (2012) Alloxan induced diabetes: mechanisms and effects, vol.3 (2), www.ijrpbsonline.com .
4. Ayesha Noor, et al (2008). Antidiabetic activity of Aloe vera and histopathology of organs in Streptozotocin induced diabetic rats. Current Science, Vol. 94(8): 1070-1076.
5. Bano B kaleem et.al (2005). Protective effects of Piper nigrum and Vinca rosea in alloxan induced diabetic rats. Indian J Physiol Pharmacol ; 49 (1): 65-67.
6. Chandra Sekhar Mohanty et.al (2013), Characterization of winged bean (Psophocarpus tetragonolobus (L.) DC.) based on molecular, chemical and physiological parameters, American Journal of Molecular Biology, vol 3, 187-197.
7. Chetna V. Ugale et.al (2011),Callus Induction Studies In *Psophocarpus Tetragonolobus* (L.).Volume 11, Issue 2, Article-005.
8. G R Juna Beegum1 et.al (2016), Natural Antioxidant Activity of *Boerhavia diffusa* L,International Journal of Pharmacognosy and Phytochemical Research,vol 8(1),8-13 .
9. Grandhi surendra et.al (2016), DPPH Free radical scavenging activity of Annona squamosa Linn leaf extracts, IOSR Journal of Pharmacy and Biological Sciences (IOSR-JPBS), Volume 11, Issue 5, Page 04-06 .
10. Hettiarachchy N.S et.al (1982), Nutritive value of winged bean, *Psophocarpus tetragonolobus, Nutrisyon* (Philippines), 7: 40-51.
11. Jain S, Bhatia et.al (2010), Antidiabetic activity of *Paspalum scrobiculatum* in alloxan induced diabetic rats, Journal of ethnopharmacology, vol 127(2), 325-328.
12. Joyamma john, (2017),Evaluate the hypoglycemic effect of *Vinca rosea* leaf Extracts in alloxan induced diabetic rats, International Journal of Science and Nature(I.J.S.N),vol.8(1),161-164.

13. Jyoti Sethi et.al (2004). Evaluation of Hypoglycemic and Antioxidant effect of Ocimum sanctum. Ind. J. Clin. Biochem., 19 (2), 152-155.
14. K. H. Lee et.al (2011),Evaluation of anti-inflammatory,antioxidant and antinoceceptive activities of six malaysian medisinal plants. Journal of Medicinal Plants Research, Vol. 5(23), pp. 5555-5563.
15. Kandasamy Murgesh et.al (2006), Antidiabetic, antioxidant and antihyperlipidemic status of *Helitropium zeylanicum* extract on streptozotocin-induced diabetic in rats; Bio. Pharm. Bull ,vol 29(11) 2202-2205.
16. Kar, A. et.al (2003), Comparative evaluation of hypoglycemic activity of some Indian medicinal plants in alloxan diabetic rats. J. Ethnopharmacology., 34: 105-108.
17. KorttA.A, (1980), Isolation and properties of a chymotrypsin inhibitor from winged bean seed *Psophocarpus tetragonolobus* (L).Biochemica Biophysica acta (BBA)-Protein structure.624, 237-248.
18. Koteswara Rao N, (2006), Anti-Hyperglycemic and Renal Protective activities of Andrographis paniculata roots Chloroform extract. Iranian J. Pharmacol. Thera., 5: 47-50.
19. L.Yoga latha et.al (2007), Antibacterial Activity and Toxicity of *Psophocarpus tetragonolobus*, Journal of Pharmaceutical biology. Volume 45, page 31-36.
20. N.Subhashini et.al(2011),Antioxidant Activity Of Trigonella Foenum Graecum Using Various In Vitro And Ex Vivo Models, International Journal of Pharmacy and Pharmaceutical Sciences , Vol 3, Issue 2,Page 96-102.
21. Neelesh Malviya et.al (2010), Antidiabetic Potential Of Medicinal Plants, Acta Poloniae Pharmaceutical and Drug Research, Vol. 67 No. 2 Pp. 113-118.
22. Oliva c. Ruma, (2016), Phytochemical screening of selected indigenous edible plants from the towns of isabela, Philippines, Asian Journal of Natural & Applied Sciences, Vol. 5(1), page 36.
23. Ozougwu et.al (2010), Studies On The Anti-Diabetic Activity Of Allium Sativum (Garlic) Aqueous Extracts On Alloxan Induced Diabetic Albino Rats, Pharmacologyonline 2: 1079-1088.
24. P Sivakumar et.al (2010),Antitumor and Antioxidant Activities of Triumfetta rhomboidea against Ehrlich Ascites Carcinoma bearing Swiss Albino mice, Research Journal of Pharmaceutical, Biological and Chemical Sciences, vol 1(4),page 486-94.

25. Prasant Kumar Rout et.al (2015), Physicochemical analysis of *Psophocarpus tetragonolobus* (L.) DC seeds with fatty acids and total lipids compositions, Journal of food science and technology, vol 52(6): 3660–3670.
26. Raghavendra H. Lakshmanashetty et.al (2010) , In vitro Antioxidant Activity of Vitex negundo L. Leaf Extracts, Chiang Mai J. Sci, vol 37(3) : 489-497.
27. Renuka C.1 et.al (2009), Evaluation of the antidiabetic effect of Trigonella foenum-graecum seed powder on alloxan induced diabetic albino rats, International Journal of PharmTech Research CODEN (USA): IJPRIF ,Vol.1(4), pp 1580-1584.
28. S.Sasidharan et.al (2008), Fungicidal Effect and Oral Acute Toxicity of *Psophocarpus tetragonolobus* Root Extract,Journal of Pharmaceutical biology.Volume 46, Page 261-265.
29. Sangameswaran. B et.al (2009), Anti-diabetic, antii-hyperlipidemic and spermatogenic effects of *Amaranthus spinosus* on streptozotocin-induced diabetic rats; Journal of Natural Medicine, 62: 79-82.
30. Shefali J. Mehta et.al (2011), Compendial Testing Method On Herbal Crude Drug - A Review, Asian J.Pharm. Res., Vol 1, Issue 2, 49-52.
31. Shokeen P et.al (2008), Antidiabetic activity of 50% ethanolic extracts of Ricinus communis and its purified fractions, Food Chem Toxicology. vol,46(11),3458-66
Sivalingam Get.al (2013) ,Anti-diabetic activity of *ribes nigrum* fruit extract in alloxan induced diabetic rats. IJPSR, ; vol. 4(3): 1196-1201.
32. Soma.S, et.al (1990).Oral hypoglycemic effect of Salacia reticulate in streptozotocin induced diabetic rats. Phytotherapy Research; 4(5): 205-206.
33. Somani R et.al **(**2005), Antidiabetic potential of Butea monosperma in rats**,** Fitoterapia., vol77(2):86-90.
34. Sonoe Ochii Yanagi Nobuko Yoshida et.al (2014), Generality and Diversity of Winged Bean (*Psophocarpus tetragonolobus*) Protein in Eight Various Line. Journal of agricultural and biological chemistry, volume 47(10), Pages 2267-227.
35. SP Anand, et.al (2018), Evaluation of antioxidant activity of some wild edible fruits collected from Boda and Kolli hills,The Journal of Phytopharmacology, vol7(2): 127-133
36. Umamaheswari S et al (2004), Hepatoprotective activity of Kadukkai Maathirai (A Siddha Polyherbal Formulation) against carbon tetrachloride induced liver damage in rats, research journal of pharmaceutical sciences; 1(4), 17-21.

37. Venkatesh S et.al (2003), Antihyperglycemic activity of Caralluma attenuata. Fitoterapia; Vol. 74, Issue 3: page no- 274-279.
38. Verma, R.S et.al (2010), Essential Oil Composition of Menthol Mint (Mentha arvensis) and Peppermint (Mentha piperita) Cultivars at Different Stages of Plant Growth from Kumaon Region of Western Himalaya. Open Access Journal of Medicinal and Aromatic Plants Vol. 1(1): 13-18.Vol.2, No.1, pp 305-309,
39. **ZHU Xiaodi LIAN Pengli et.al (2015),**Antioxidant activities of Psophocarpus tetragonolobus(L.),Journal of henan university(medical science),Available from http://en.cnki.com.cn/Article_en/CJFDTOTAL-KFYZ201501005.html.